INDEX

Adalia bipunctata, 487, 489, 490, 491.
Agrilus viridis, 134.
Agromyza jucunda, 431.
Agrotis jaculifera, 394.
Air conditioning apparatus, 107–111.
Alabama argillacea, 429.
Aldrich, J. M., 151, 242.
Alfalfa injured by thrips, 360.
  looper, 286–291.
Allorhina nitida parasite, 151.
American Association of Economic Entomologists:
  Apiary inspection, report of section, 119–128.
  Committees, ix.
    Bibliography, 25.
    Executive, 5.
    Membership, 23, 27.
    Resolutions, 24, 499.
  Constitution, xxi.
  Horticultural Inspectors Proceedings, 129–135, 268–276.
  Meetings, list, x.
  Members, list, xii.
  Membership committee, 23, 27.
  Officers, ix.
  Resolution committee, 24, 499.
  Secretary, report, 3.
  Special committee, 499.
  Special meeting, report, 441–500.
  Twenty-seventh meeting, proceedings, 1–135, 161–292, 317–338.
American economic entomology, bibliography, 25.
Ammonia, 94.
Anagrus frequens, 454.
Angoumois grain moth, 233.
Anthrax, 328.
Anthrenus verbasci, 430.
Antonina crawi, 551.
Aphids and fire blight, 402–403, 551.
Apiary inspection, report of section, 119–128; record system, 121–123.
Apicultural inspection, 123–128.
Apple bark disease and snowy tree-cricket, 535–541.
  flea weevil, 279–286.
  leaf hopper, black, 415–419.
  maggot, 311.
  red bug, 293–295.
    false, 295, 296.
Arsenate of lead, 459, 466; toxic values, 481–486.
Arsenate of lime, 194–197.
Arsenical poisoning, 226.
  residues on fruit and grass, 191.
Army worm, 220–222, 236; parasites, 299–302.
Asteromyia leviana, 406.
Autographa gamma californica, 286–291.
Bacillus amylovorus, 402–403.
Back, E. A., 475.
Ball, E. D., 475, 496, 525.
Barium sulphide, 206.
Barred-winged onion fly, 349.
Bean, Ceratoma trifurcata injuring, 261.
Bean leaf weevil, 261–266.
Bees and poisoned bran mash, 234.
Beet leaf hopper, 496.
Beetle, flea, catcher, 240–241.
Beneficial insects, introduction in Hawaiian Islands, 450–456.
Bipalium kewense, 404.
Bishopp, F. C., 54, 317, 466.
Black cricket, 223.
  onion fly, 348.
Blakeslee, E. B., 103.
Borden, A. D., 354.
Bourne, A. I., 276.
Box leaf miner, 94–95, 306.
Bragg, L. C., 97.
Bran mash, poisoned, 219–223.
Braucher, R. W., 429.
Britton, W. E., 72, 121, 379.
Brown lace-wing, 350–354.
Brown-tail moth, 85–88.
Brues, C. T., 382.
Burgess, A. F., 307, 496.
Burrill, A. C., 552, 553.

Cabbage curculio, 429.
Cabbage flea-beetle, western, 429.
Calcium arsenate, 194–197.
Calliphora dux, 320.
  erythrocephala, 327.
  oceaniæ, 320.
  rufifacies, 320.
  villosa, 320.
Callopistria floridensis, 134.
Camnula pellucida, 526, 528, 530.
Capsids, observations on, 293–298.
Carbon bisulfid, 94, 233.
Carbon tetrachloride, 94.
Carpophilus hemipterus, 396–400.
Celatoria diabroticæ, 264.
Ceratitis capitata, 329–338.
Ceratoma trifurcata, 261–266.
Ceromasia sphenophori, 455.
Ceutorhynchus rapæ, 429.
Chalcis obscurata, 452.
Chapman, J. W., 140.
Citricola scale, 291–292.
Citrus trees, spraying scheme, 161–164.
Chætopsis ænea, 349.
Chermes cooleyi, 97.
Chicken flea, 492–495.
Chickens and poisoned bran marsh, 235.
  killed by rose chafers, 547.
Childs, Leroy, 457.
Chinch bug egg parasite, 248–261.

INDEX

Adalia bipunctata, 487, 489, 490, 491.
Agrilus viridis, 134.
Agromyza jucunda, 431.
Agrotis jaculifera, 394.
Air conditioning apparatus, 107–111.
Alabama argillacea, 429.
Aldrich, J. M., 151, 242.
Alfalfa injured by thrips, 360.
  looper, 286–291.
Allorhina nitida parasite, 151.
American Association of Economic Entomologists:
  Apiary inspection, report of section, 119–128.
  Committees, ix.
    Bibliography, 25.
    Executive, 5.
    Membership, 23, 27.
    Resolutions, 24, 499.
  Constitution, xxi.
  Horticultural Inspectors Proceedings, 129–135, 268–276.
  Meetings, list, x.
  Members, list, xii.
  Membership committee, 23, 27.
  Officers, ix.
  Resolution committee, 24, 499.
  Secretary, report, 3.
  Special committee, 499.
  Special meeting, report, 441–500.
  Twenty-seventh meeting, proceedings, 1–135, 161–292, 317–338.
American economic entomology, bibliography, 25.
Ammonia, 94.
Anagrus frequens, 454.
Angoumois grain moth, 233.
Anthrax, 328.
Anthrenus verbasci, 430.
Antonina crawi, 551.
Aphids and fire blight, 402–403, 551.
Apiary inspection, report of section, 119–128; record system, 121–123.
Apicultural inspection, 123–128.
Apple bark disease and snowy tree-cricket, 535–541.
  flea weevil, 279–286.
  leaf hopper, black, 415–419.
  maggot, 311.
  red bug, 293–295.
    false, 295, 296.
Arsenate of lead, 459, 466; toxic values, 481–486.
Arsenate of lime, 194–197.
Arsenical poisoning, 226.
  residues on fruit and grass, 191.
Army worm, 220–222, 236; parasites, 299–302.
Asteromyia leviana, 406.
Autographa gamma californica, 286–291.
Bacillus amylovorus, 402–403.
Back, E. A., 475.
Ball, E. D., 475, 496, 525.
Barium sulphide, 206.
Barred-winged onion fly, 349.
Bean, Ceratoma trifurcata injuring, 261.
Bean leaf weevil, 261–266.
Bees and poisoned bran mash, 234.
Beet leaf hopper, 496.
Beetle, flea, catcher, 240–241.
Beneficial insects, introduction in Hawaiian Islands, 450–456.
Bipalium kewense, 404.
Bishopp, F. C., 54, 317, 466.
Black cricket, 223.
  onion fly, 348.
Blakeslee, E. B., 103.
Borden, A. D., 354.
Bourne, A. I., 276.
Box leaf miner, 94–95, 306.
Bragg, L. C., 97.
Bran mash, poisoned, 219–223.
Braucher, R. W., 429.
Britton, W. E., 72, 121, 379.
Brown lace-wing, 350–354.
Brown-tail moth, 85–88.
Brues, C. T., 382.
Burgess, A. F., 307, 496.
Burrill, A. C., 552, 553.

Cabbage curculio, 429.
Cabbage flea-beetle, western, 429.
Calcium arsenate, 194–197.
Calliphora dux, 320.
  erythrocephala, 327.
  oceaniæ, 320.
  rufifacies, 320.
  villosa, 320.
Callopistria floridensis, 134.
Camnula pellucida, 526, 528, 530.
Capsids, observations on, 293–298.
Carbon bisulfid, 94, 233.
Carbon tetrachloride, 94.
Carpophilus hemipterus, 396–400.
Celatoria diabroticæ, 264.
Ceratitis capitata, 329–338.
Ceratoma trifurcata, 261–266.
Ceromasia sphenophori, 455.
Ceutorhynchus rapæ, 429.
Chalcis obscurata, 452.
Chapman, J. W., 140.
Citricola scale, 291–292.
Citrus trees, spraying scheme, 161–164.
Chætopsis ænea, 349.
Chermes cooleyi, 97.
Chicken flea, 492–495.
Chickens and poisoned bran marsh, 235.
  killed by rose chafers, 547.
Childs, Leroy, 457.
Chinch bug egg parasite, 248–261.

Chlorochroa uhleri, 308.
Chorizagrotis agrestis, 443.
Chrysanthemum midge, 267.
Chrysanthemum, Rhopalomyia hyphogæa injuring, 267.
Chysomyia macellaria, 320–322, 323, 324, 325–326, 469.
Chrysopa rufilabris, 431.
Clausen, C. P., 487.
Clean town contest, 175–177.
Coccinella californica, 487, 489, 490, 491.
trifasciata, 487, 489, 490, 491.
Coccinellidæ, aphid feeding, 487–491.
Coccus citricola, 291–292.
Cockerell, T. D. A., 549.
Codling moth, spraying methods, 164–167.
Colorado aphids, 97–103.
Conchuela, 232.
Contarinia sorghicola, 230–232.
Cooley, R. A., 47, 442.
Corn-ear worm, 211–216.
Corn silk beetle, 507.
Cory, E. N., 186, 270.
Cotton moth, 429.
seed fumigation, 400–402.
seed oil, 298.
worm, 192.
Cowpeas, Ceratoma trifurcata injuring, 261.
Craighead, F. C., 513.
Cranberry root worm, 193.
Criddle mixture, 533–534.
Crioceris asparagi, 549.
Crosby, C. R., 409, 541.
Crude oil, 461–464.
Crumb, S. E., 392.
Cryptolæmus montrouzieri, 452.
Cryptorhynchus lapathi, 522–525.
Cucumber beetle, western striped, 515, 519–520.
beetle, western twelve-spotted, 515–519.
Cutworms, 349; tobacco, key to, 392–395.
Cutworm, variegated, 222–223.
Cycloneda abdominalis, 487, 489, 490, 491.
sanguinea, 487.
Cyllene pictus, 513, 549.
robiniae, 513.
Cynomyia cadaverina, 327.

Damaster blaptoides, 135.
Dasyneura torontoensis, 405.
Davidson, W. M., 419.
Davis, J. J., 135.
Dean, G. A., 107, 219, 429.
Dermacentor venustus, 47–53.
Dermatobia hominis, 319.
Diabrotica soror, 515–519.
trivittata, 519–520.
Diachasma tryoni, 456.
Diatræa saccharalis, 496–499.
Dinoderus substriatus, 388.
Diprion simile, 379–382.
Douglass, B. W., 431.
Dove, W. E., 54.
Dræculacephala reticulata, 430.
Dried fruit beetle, 396–400.
Dust sprays for corn-ear worm, 211–214.

Echidnophaga gallinacea, 492–495.
Edibility of insects, 549.
Ellis, W. O., 520.
Economic Entomologists, Pacific Slope Association, 449.
entomology, college extension work in, 442–449.
entomology, notes on progress, 113–118.
entomology, some present needs, 30–40.
results of the year, 234–238.
Enchenopa binotata, 368–371.
English sparrow, 553.
Entomological investigations, report on, 8–20.
museum, University of Kansas, 305.
Entomologists, academic training, 72–78.
Eriococcus borealis, 549.
Essig, E. O., 396.
Eucactophagus graphipterus, 133.
Eucallipterus flavus, 423.
Euceraphis gillettei, 421.
Eudiagogus rosenschoeldi, 551.
Eumicrosoma benefica, 248–261.
European mole cricket, 500–501.
Euthrips occidentalis, 360.
tritici, 359, 360.
Euxesta notata, 349.
Euxoa messoria, 394.

Felt, E. P., 94, 149, 160, 227, 267, 308, 405, 430, 550.
Feltia annexa, 394.
gladiaria, 394.
jaculifera, 394.
malefida, 394.
Feltiella davisi, 406.
Fernald, H. T., 30.
Fire blight and aphids, 402–403, 551.
Fletcher, T. B., 154.
Flies causing myiasis, 317–329.
Foreign plant pests, manual, 430.
Foul brood distribution, 121.
Fowls and grasshoppers, 226.
Frauds, semi-frauds and questionables, 199.
Fruit-fly, Mediterranean, 475.
Fruit tree leaf-roller, spraying notes on, 457.
Fulton, B. B., 535.
Fumigation, relative size in, 302–304.
Funkhouser, W. D., 368.

Gall midges, 405–409, 550.
Garman, H., 403.

Gates, B. N., 123.
Gillette, C. P., 97, 375.
Gipsy moth conference, 306–307; work, 496.
Glaser, R. W., 140.
Glenn, P. A., 276.
Glischrochilus fasciatus, 350.
Gloyer, W. O., 535.
Gonatocerus ovicenatus, 545.
Grasshopper outbreaks, 475.
Grasshoppers, 219–222, 227–229, 527–535; estimating the number of, 525–527.
Gray, G. P., 487.
Gryllotalpa gryllotalpa, 500–501.
Gryllus pennsylvanicus, 223, 308.

Habitat, original of insect pests, 171–174.
Hadley, C. H., 88, 409.
Handbook of medical entomology, 310.
Harned, R. W., 508.
Haseman, L., 192, 214.
Headlee, T. J., 40, 271.
Heliophila multilinea, 389–392.
subpunctata, 389–392.
unipuncta, 389–392; parasites, 299–302. See also Leucania.
Heliothis obsoleta, 211–216.
Hemerobius pacificus, 350–354.
Herrick, G. W., 180, 313.
Heterocordylus malinus, 293–295.
Heterodera schachtii, 404.
Hewitt, C. Gordon, 85, 310.
Hinds, W. E., 400.
Hippodamia ambigua, 487, 489, 490, 491.
convergens, 487, 489, 490, 491.
Hopperdozer, 531–533.
Horticultural Inspectors Proceedings, 129–135.
Host labels for parasites, 550.
House-fly, breeding material, 64.
biology, 54–71.
copulation, 55.
developmental periods, 60.
hibernation, 65.
its structure, habits, development, relation to disease and control, 310.
Howard, L. O., 113, 549, 550.
Hunter, S. J., 234.
Hypoderma bovis, 318.
lineata, 318.

Idiocerus alternatus, 541.
gemmisimulans, 542–544, 545.
provancheri, 415–419.
Ichneumon lætus, 514–515.
Illingworth, J. F., 492.
Insect control, essentials, 271–276.
edibility 549.
importations, recent in New Jersey, 133–135.
pests, original habitat, 171–174.
Insects collected on nursery stock, 268–270.
North American, key to families, 382.
of economic importance, outlines in economic entomology, 313.
Intermittent, seasonal spraying, 236.
Iridomyrmex humilis, 133.

Johanssen, O. A., 310.
Juniper plant bug, 308.

Kerosene emulsion, 190.
emulsion and sodium arsenate, 513.
traps, 329–338.
Keys to the families of North American Insects, 382.
Knight, H. H., 293, 514.

Laake, E. W., 466.
Lamson, G. H., 547.
Larrimer, W. H., 430.
Leaf-roller, fruit tree, 180–184.
Lecanium mangiferæ, 388.
Leonard, M. D., 361, 415, 541.
Leptobyrsa explanata, 409–414.
Lepidosaphes ulmi, 371–375.
Leptinotarsa decemlineata, 520–521.
Leucania unipuncta, 220–222. See also Heliophila.
Leucaspis japonica, 160.
Life zones, 36, 90.
Lucilia cæsar, 467.
sericata, 326, 467, 469, 472–473.
Luperodes varicornis, 507.
Lygidea mendax, 295–296.
Lygus invitus, 296–297.
pratensis, 361–367.

Macrodactylus subspinosus, 547–548.
Macrodyctium omiodivorum, 452.
Macrosiphum heucheræ, 427.
Malacosoma pluvialis, 481.
Mamestra legitima, 393.
meditata, 393.
renigera, 393.
Matheson, Robert, 522.
McColloch, J. W., 211.
McConnell, W. R., 551.
Mediterranean fruit-fly, 329–338, 475.
Melander, A. L., 382, 475.
Melanoplus atlanis, 528.
bivittatus, 528.
destructor, 530.
differentialis, 530.
minor, 528.
uniformis, 530.
Merrill, J. H., 402.
Mesogramma polita, 338–342.
Metcalf, C. L., 240.
Miscible oil, 462–464.
Monarthropalpus buxi, 94–95.
Monell, J. T., 503.
Mosquito control in a limited locality, 40–47.
Moznette, G. F., 350.

Mycodiplosis fungiperda, 407.
macgregori, 149.
Myelophilus piniperda, 134.
Myiasis, flies causing, 317–329.
Myzocallis pasaniae, 424.
Myzus braggii, 375–379.

New contact insecticide, 206–210.
Newell, Wilmon, 120, 230.
Nocotine sulphate-bordeaux combination, 199–203.
Noctua badinodis, 394.
clandestina, 394.
c-nigrum, 393.

Oberea tripunctata, 79–85.
ulmicola, 79.
Œcanthus niveus, 535–541.
Œstrus ovis, 318.
Oil, crude, 461–464.
miscible, 462–464.
O'Kane, W. C., 191, 311.
Olla abdominalis, 487, 489, 490, 491.
oculata, 487, 489, 490, 491.
Onion fly, imported, 342–348.
maggot, 276–279.
thrips, 349.
Opius humilis, 388, 456.
Orchestes canus, 276–286.
Organization in agricultural college extension work in economic entomology, 442–449.
Otiorhynchus sulcatus, 552.
Ox warble, 318.
Oxybelus 4-notatus, 344.
Oyster shell scale, 371–375.

Paracalocoris colon, 297.
Paraffin-oil emulsion, 162.
Parallelodiplosis corticis, 407.
Paranagrus obtabilis, 454.
perforator, 454.
Parker, J. R., 286.
Parker, R. R., 246.
Parman, D. C., 54.
Parrott, P. J., 164, 204, 535.
Peach borer, banding materials for, 106–107; mechanical protector for, 103–105.
Pemberton, C. E., 475.
Pemphigus betæ, 97.
Pennington, W. E., 433.
Pentatoma ligata, 232.
Peridroma incivis, 393.
saucia, 393, 394.
Perkinsiella saccharicida, 454.
Phillips, E. F., 121.
Phorbia cepetorum, 342–348.
Phormia regina, 322, 327, 469–472.
Phyllotreta pusilla, 429.
vittata, 429.
Phylloxera popularia, 420.
salicola, 419.
Pine sawfly, European, 379–382.
tar creosote, 189.
Plant lice, western, 419–429.
Poisoned baits, 528–530, 533–535.
bait spray for onion fly, 342–348.
bran mash, 219–223; and bees, 234; chickens, 235.
Pomphopœa ænea, 431.
Poplar and willow borer, 522–525.
Porthetria dispar, 134.
Presidential address, 30–40; discussion, 89–92.
Prionellus monilis, 405.
Prociphilus fraxini-dipetalæ, 421.
Prodenia commelinæ, 393.
ornithogalli, 393.
Pseudococcus kraunhiae, 552.
Pteronus hudsonii, 552.

Quayle, H. J., 291.

Rabbit injury, prevention, 270.
Ravinia peniculata, 247.
Retinidiplosis palustris, 408.
Rhabdophaga aceris, 549.
Rhododendron lace-bug, 409–414.
Rhogas terminalis, 433.
Rhopalomyia hypogæa, 267.
Rhopalosiphum hippohæs, 375–379.
Richardson, C. H., 338.
Riley, W. A., 310.
Robinson, Elizabeth, 549.
Romalea microptera, 529.
Rose chafer, 547–548.
Ruggles, A. G., 79.
Russell, H. M., 433.

Safro, V. I., 199.
San José scale parasites, 238; susceptibility, 475–480; and birds, 553.
Sarcophaga lambens, 319.
pyophila, 319.
utilis, 151.
Sarcophagidæ, economic relations, 242–246.
Sasscer, E. R., 268.
Scammell, H. B., 193.
Schizoneura americana, 99.
cratægi, 100.
lanigera, 100.
rileyi, 99.
Schœne, W. J., 204.
Scott, W. M., 194, 206.
Screw worm, 322.
Sell, R. A., 515.
Severin, H. C., 329, 342.
Severin, H. H. P., 329, 342.
Sheep bot, 318.
Shelford, V. E., 171.
Sherman, Jr., Franklin, 299.
Sitotroga cerealella, 233.
Snowy tree-cricket, 535–541.
Sodium arsenate-kerosene emulsion, 513.
Soft maple leaf midge, 549.
Sorghum midge, 230–232.
South Indian insects, 154.
Spotted fever tick, 47–53.

Spraying notes on fruit tree leaf-roller, 457.
Stable fly, 350.
Stomoxys calcitrans, 350.
Striped turnip flea-beetle, 429.
Sudan grass enemies, 230–234.
Sulfur as an insecticide, 487.
Sulphides and polysulphides, insecticidal properties, 204–205.
Surface, H. A., 199, 238.
Swezey, O. H., 450.
Syrphus fly, corn-feeding, 338–342.

Tarnished plant bug, 361–367; false, 296–297.
Tarter, H. V., 481.
Termes lucifugus, 308.
Thecabius populicaulis, 420.
populiconduplifolius, 99.
Thrips tabaci, 349.
Thysanopters, mouthparts and injuries by, 354–360.
Titus, E. G., 175.
Tower, D. G., 431.
Toxicity, theory of, 509–512.
Trioza magnoliae, 551.
Tritoxa flexa, 348.
Turkeys and grasshoppers, 226.

Underground insects, methods of study, 135–139.
Unique type of insect injury, 261–266.

Vickery, R. A., 389.

Walker, H. C., 306.
Water under pressure for mealy bug, 304–305.
Webster, F. M., 527.
Webster, R. L., 371.
Weiss, H. B., 133, 500, 552.
Weldon, G. P., 450.
Wilson, H. F., 481.
Wilt, list of affected insects, 140–149.
Wilt polyhedra, 146.
Wohlfahrtia magnifica, 320.
Wolcott, G. N., 496.
Woodworth, C. W., 302, 509
Woolly aphis, 186–190, 450.
Wool maggots, 466–474; control, 473–474.
Worms, introduced, 403–404.

Yothers, W. W., 161, 298, 304.